

B7-H3 Protein, Human, Recombinant (aa 29-245, His)

General Information

Synonyms:	PRO352;B7H3; UNQ309;PSEC0249;CD276;B7-H3;B7 homolog 3
Protein Construction:	Leu29-Pro245
Species:	Human
Expression Host:	HEK293 Cells
Accession:	Q5ZPR3-2
Molecular Weight:	24.7 kDa (predicted). Due to glycosylation, the protein migrates to 40-50 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Immobilized Human B7-H3, His Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Anti-B7-H3 Antibody, hFc Tag with the EC50 of 29.4ng/ml determined by ELISA.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 µg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

B7-H3, a member of the B7 family of immunomodulatory molecules, is overexpressed in a wide range of solid cancers. B7-H3 binds to activated T cells via an as yet unidentified receptor. In assays using sub-optimal amount so anti-CD3 stimulation, 2Ig-B7-H3 enhances T cell proliferation, T cell interferon-gamma (IFN-gamma) production, and cytotoxic T cells induction.

Reference

Fodstad O, et al. The Immunoregulatory Protein Human B7H3 is a Tumor-Associated Antigen that Regulates Tumor Cell Migration and Invasion[J]. Current Cancer Drug Targets, 2008, 8(5)

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481