

BSP11 Protein, Mouse, Recombinant (His)

General Information

Synonyms:	IBSP;SP 11;BSP;BSP 11;BNSP;BSP-11;BSP 2;SP11;SP-11
Protein Construction:	Phe17-Gln324
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	Q61711
Molecular Weight:	34.9 kDa (predicted). Due to glycosylation, the protein migrates to 68-80 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/ μ g of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μ m filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μ g/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Osteopontin (OPN), bone sialoprotein (BSP11), and osteonectin (ON) belong to a family of glycoproteins, which have been linked to cancer metastasis and progression. Here, we report on the selection of antisense oligonucleotides (ASOs), which are effective in reducing their protein levels.

Reference

Adwan H, Bäuerle TJ, Berger MR. Downregulation of osteopontin and bone sialoprotein II is related to reduced colony formation and metastasis formation of MDA-MB-231 human breast cancer cells. *Cancer Gene Ther.* 2004 Feb;11(2):109-20. doi: 10.1038/sj.cgt.7700659. PMID: 14647232.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481