

LILRA3/CD85e Protein, Human, Recombinant (His & Avi)

General Information

Synonyms:	HM43;LIR-4HM43e3;HM31;ILT-6;LIR4;CD85e;LILRA3;ILT6
Protein Construction:	Gly24-Glu439
Species:	Human
Expression Host:	HEK293 Cells
Accession:	Q8N6C8
Molecular Weight:	47.9 kDa (predicted). Due to glycosylation, the protein migrates to 68-72 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Human LILRA3, also known as Leukocyte Ig-like receptor 4 (LIR-4), CD85 antigen-like family member E, Immunoglobulin-like transcript 6 (ILT-6) and monocyte inhibitory receptor HM43/HM31, is a 70 kD Ig superfamily member that belongs to the leukocyte receptor complex/cluster. Mature LILRA3 is 416 amino acids (aa) in length and contains four C2-type Ig-like domains. Unlike other LILR family members, LILRA3 is actively secreted. LILRA3 acts as soluble receptor for class I MHC antigens. LILRA3 binds both classical and non-classical HLA class I

A DRUG SCREENING EXPERT

molecules but with reduced affinities compared to LILRB1 or LILRB2. Binds with high affinity to the surface of monocytes, leading to abolish LPS-induced TNF-alpha production by monocytes.

Reference

Ordóñez D, et al. Multiple sclerosis associates with LILRA3 deletion in Spanish patients[J]. Genes & Immunity, 2009, 10(6):579-585.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481