

IL-31RA Protein, Human, Recombinant (aa 20-519, His)

General Information

Synonyms:	GLM-RMGC125346;CRL3;GLM-R;PRO21384;Gp130-like receptor;CRL3Glmr;IL-31R-alpha;IL-31RA;GPL;IL-31R- α ;ZcytoR17
Protein Construction:	Ala20-Glu519
Species:	Human
Expression Host:	HEK293 Cells
Accession:	Q8NI17-1
Molecular Weight:	58.2 kDa (predicted). Due to glycosylation, the protein migrates to 90-116 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	<ol style="list-style-type: none">1. Immobilized Human IL-31 RA, His Tag at 0.5μg/ml (100μl/well) on the plate. Dose response curve for Anti-IL-31 RA Antibody, hFc Tag with the EC50 of 6.5ng/ml determined by ELISA (QC Test).2. Human IL-31, His Tag immobilized on CM5 Chip can bind Human IL-31 RA, His Tag with an affinity constant of 82.4nM as determined in a SPR assay.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/ μ g of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μ m filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:	Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μ g/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.
Stability & Storage:	<p>It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.</p> <p><small>Actual storage temperature shall be subject to the COA.</small></p>
Shipping:	In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Interleukin-31 receptor A (IL-31RA) is a newly identified type I cytokine receptor, that is related to gp130, the

common receptor of the interleukin (IL) -6 family cytokines. IL-31RA forms a functional receptor complex for IL-31 together with the beta subunit of oncostatin M receptor (OSMRbeta). OSMRbeta is expressed in a subset of small-sized nociceptive neurons of adult dorsal root ganglia (DRGs).IL-31 and OSM may thus have redundant functions in the development of OSMRbeta-expressing neurons.

Reference

Bando T, et al. Complete overlap of interleukin-31 receptor A and oncostatin M receptor beta in the adult dorsal root ganglia with distinct developmental expression patterns. *Neuroscience*. 2006 Nov 3;142(4):1263-7doi: 10.1016/j.neuroscience.2006.07.009. Epub 2006 Aug 22. PMID: 16926070.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481