

CD3 epsilon/CD3e Protein (Primary Amine Labeling), Cynomolgus, Recombinant (His),

General Information

Synonyms:	TCRE;CD3e;CD3-epsilon;FLJ18683;CD3-ε;CD3E/CD3 ε;IMD18;T3E
Protein Construction:	Gln22-Asp117
Species:	Cynomolgus
Expression Host:	HEK293 Cells
Accession:	Q95LI5
Molecular Weight:	11.7 kDa (predicted). Due to glycosylation, the protein migrates to 13-16 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Immobilized Anti-CD3 Antibody, hFc Tag at 0.5µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Cynomolgus CD3E, His Tag with the EC50 of 0.87ng/ml determined by ELISA.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:
Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 µg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

CD3E, is a single-pass type I membrane protein. CD3 (cluster of differentiation 3) T cell co-receptor helps to activate both the cytotoxic T cell (CD8 naive T cells) and also T helper cells (CD4 naive T cells). It consists of a protein complex and is composed of four distinct chains. In mammals, the complex contains a CD3γ chain, a CD3δ chain,

and two CD3ε chains.

Reference

Kroesen B J, et al. Approaches to lung cancer treatment using the CD3ε×GP-2-directed Bispecific Monoclonal Antibody BIS-1[J]. Cancer Immunology Immunotherapy, 1997, 45(3-4):203-206.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481