

Siglec-8 Protein, Human, Recombinant (His & Avi), Biotinylated

General Information

Synonyms:	SIGLEC8;CDw329;SAF2SAF-2;SAF2;SIGLEC8L;Siglec-8;MGC59785
Protein Construction:	Met17-Ala363
Species:	Human
Expression Host:	HEK293 Cells
Accession:	Q9NYZ4
Molecular Weight:	40.7 kDa (predicted). Due to glycosylation, the protein migrates to 50-60 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 95% as determined by Tris-Bis PAGE
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 µg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Siglec-8, also known as SAF, is an approximately 75 kDa transmembrane glycoprotein in the Siglec family of sialic acid-binding immune regulatory molecules. Mature human Siglec-8 consists of a 347 amino acid (aa) extracellular domain (ECD) with three Ig-like domains. Putative adhesion molecule that mediates sialic-acid dependent binding to red blood cells. Preferentially binds to alpha-2,3-linked sialic acid. Also binds to alpha-2,6-linked sialic acid.

Reference

Macauley M S, et al. Siglec-mediated regulation of immune cell function in disease[J]. Nature Reviews Immunology, 2014, 14(10):653-666.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481