

CD161 Protein, Cynomolgus, Recombinant (His & Avi)

General Information

Synonyms:	HNKR-P1a;NKR P1;KLRB1;Ly59;NKR P1ANKR;NKR-P1;CLEC5B;NKR;CD161;NKR P1A;NKR-P1A
Protein Construction:	Gln67-Leu227
Species:	Cynomolgus
Expression Host:	HEK293 Cells
Accession:	XP_005570142.1
Molecular Weight:	21.5 kDa (predicted). Due to glycosylation, the protein migrates to 37-42 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Immobilized Cynomolgus CD161, His Tag at 2µg/ml (100µl/Well) on the plate. Dose response curve for Anti-CD161 Antibody, hFc Tag with the EC50 of 35.0ng/ml determined by ELISA.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 µg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

CD161 (NKR P1) is a lectin-like receptor present on NK cells and rare T-cell subsets. We have observed CD161 expression in some cases of T-cell prolymphocytic leukemia (T-PLL) and found it to be useful in follow-up and detection of disease after treatment.

Reference

Gilles SR, et al. CD161 Is Expressed in a Subset of T-Cell Prolymphocytic Leukemia Cases and Is Useful for Disease Follow-up. Am J Clin Pathol. 2019 Sep 9;152(4):471-478. doi: 10.1093/ajcp/aqz060. PMID: 31305900.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481