

## Bat coronavirus HKU9 Spike glycoprotein (His)

## General Information

Synonyms:	S glycoprotein;E2;Spike glycoprotein;Peplomer protein
Protein Construction:	341-533 aa
Species:	BtCoV
Expression Host:	E. coli
Accession:	A3EXG6
Molecular Weight:	27.6 kDa (predicted)
AA Sequence:	RAQVAGFVRVTQRGSYCTPPYSVLQDPPQPVVWRRYMLYDCVFDFTVVVDSLPTHQLQCYGVSPRRLASMC YGSVTLDVMRINETHLNNLNFNRVPDFTSLYNYALPDNFYGLHAFYLNSTAPYAVANRFPKPGGRQSNSAFI DTVINAAHYSPFSYVYGLAVITLKPAAGSKLVCPVANDTVVITDRVCQY

## QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

## Preparation and Storage

Reconstitution:	Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.
Stability & Storage:	Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots. <small>Actual storage temperature shall be subject to the COA.</small>
Shipping:	In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

## Protein Background

attaches the virion to the cell membrane by interacting with host receptor, initiating the infection.; mediates fusion of the virion and cellular membranes by acting as a class I viral fusion protein. Under the current model, the

## A DRUG SCREENING EXPERT

---

protein has at least three conformational states: pre-fusion native state, pre-hairpin intermediate state, and post-fusion hairpin state. During viral and target cell membrane fusion, the coiled coil regions (heptad repeats) assume a trimer-of-hairpins structure, positioning the fusion peptide in close proximity to the C-terminal region of the ectodomain. The formation of this structure appears to drive apposition and subsequent fusion of viral and target cell membranes.; Acts as a viral fusion peptide which is unmasked following S2 cleavage occurring upon virus endocytosis.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481