

Bovine coronavirus (strain OK-0514) Spike glycoprotein (His)

General Information

Synonyms: S glycoprotein;E2;Peplomer protein;Spike glycoprotein

Protein Construction: 314-634 aa

Species: BCoV

Expression Host: HEK293 Cells

Accession: Q9QAQ8

Molecular Weight: 37.2 kDa (predicted)

AA Sequence: TVQPIADVRRIPNLPDCNIEAWLNDKSVPSPLNWERKTFNSCNFNMSSLMSFIQADSFTCNNIDA AKIYGMCFSSITIDKFAIPNGR KVDLQLGNLGYLQSFNYRIDTTATSCQLYYNLP AANVSVSRFNPSIWNRRFGFTEQSVFKPQPAGVFTDHDVVYAQHCFKAPT NFCPCCKLDGSLCVGSGSIDAGYKNTGIGTCPAGTNYLTCHNAAQCGCLCTPDPITSKATGPYKCPQTKYLVGIGEHCSGLAIKSDYCGGNPCSCRPAFLGWSVDSCLQGDR CNIFANFI LHDVNSGTTCCSTDLQKSNTDIILGVCVNY

QC Testing

Biological Activity: Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.

Purity: > 85% as determined by SDS-PAGE.

Endotoxin: < 1.0 EU/μg of the protein as determined by the LAL method.

Formulation: If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

attaches the virion to the cell membrane by interacting with host receptor, initiating the infection.; mediates fusion of the virion and cellular membranes by acting as a class I viral fusion protein. Under the current model, the protein has at least three conformational states: pre-fusion native state, pre-hairpin intermediate state, and post-fusion hairpin state. During viral and target cell membrane fusion, the coiled coil regions (heptad repeats) assume a trimer-of-hairpins structure, positioning the fusion peptide in close proximity to the C-terminal region of the ectodomain. The formation of this structure appears to drive apposition and subsequent fusion of viral and target cell membranes.; Acts as a viral fusion peptide which is unmasked following S2 cleavage occurring upon virus endocytosis.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481