

Adrenodoxin reductase Protein, Bovine, Recombinant (E. coli, His)

General Information

Synonyms:	NADPH:adrenodoxin oxidoreductase, mitochondrial;Adrenodoxin reductase;Ferredoxin--NADP(+) reductase (Ferredoxin reductase);ADXR;mitochondrial;FDXR;AR
Protein Construction:	33-492 aa
Species:	Bovine
Expression Host:	E. coli
Accession:	P08165
Molecular Weight:	55.8 kDa (predicted)
AA Sequence:	STQEQTPQICVVGSGPAGFYTAQHLLKHHSRAHVDIYEKQLVFPGLVRFVAPDHPEVKNVINTFTQTARSDR CAFYGNVEVGRDVTVQELQDAYHAVVLSYGAEDHQALDIPGEELPGVFSARAFVWYNGLPENRELAPDLS CDTAVILGQGNVALDVARILLTPPDHLEKTDITEAALGALRQSRVKTWVWVGRGRLQVAFTIKELREMIQLPGT RPMLDPADFLGLQDRIKEAARPRKRLMELLRTATEKPGVEEAARRASASRAWGLRFFRSPQQVLPSPDGRR AAGIRLAVTRLEGIGEATRAVPTGDVEDLPCGLVLSSIGYKSRPIDPSVFPDKLGVVPMMEGRVVDVPGLYCS GWVKGPTGVITTTMTDSFLTGQILLQDLKAGHLPSGPRPGSAFIKALLDSRGVWPVFSFSDWEKLDAAEEVSRG QASGKPREKLLDPQEMLRLLGH

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 85% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Tris-based buffer, 50% glycerol

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

A DRUG SCREENING EXPERT

Serves as the first electron transfer protein in all the mitochondrial P450 systems including cholesterol side chain cleavage in all steroidogenic tissues, steroid 11-beta hydroxylation in the adrenal cortex, 25-OH-vitamin D3-24 hydroxylation in the kidney, and sterol C-27 hydroxylation in the liver.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481