

DHAK Protein, Citrobacter freundii, Recombinant (His & SUMO)

General Information

Synonyms: dhaK;Dihydroxyacetone kinase;DHA kinase;Glycerone kinase

Protein Construction: 356-548 aa

Species: Citrobacter freundii

Expression Host: E. coli

Accession: P45510

Molecular Weight: 35.9 kDa (predicted)

AA Sequence: ALVAGIVELVTATLSLDLETHLNALDAKVGDDGTGSTFAAAAAREIASLLHRQQLPLNNLATLFFALIGERLTVVMG
GSSGVLMSIFFTAAGQKLEQGANVVEALNTGLAQMKFYGGADEGDRMTIDALQPALTSLLAQPKNLQAAFD
AAQAGAERTCLSSKANAGRASYLSSESLLGNMDPGAQRLAMVFKALAE

QC Testing

Biological Activity: Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.

Purity: > 90% as determined by SDS-PAGE.

Endotoxin: < 1.0 EU/μg of the protein as determined by the LAL method.

Formulation: Tris-based buffer, 50% glycerol

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Catalyzes the phosphorylation of dihydroxyacetone.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481