

Ex-FABP Protein, Coturnix coturnix japonica, Recombinant (His & Myc)

General Information

Synonyms:	Lipocalin Q83; Extracellular fatty acid-binding protein; Ex-FABP
Protein Construction:	21-178 aa
Species:	Coturnix coturnix japonica
Expression Host:	E. coli
Accession:	Q9I9P7
Molecular Weight:	25.5 kDa (predicted)
AA Sequence:	AATVPDRSEIAGKQWYVVALASNTEFFLREKDKMKMAMARISFLGEDELKVSYAVPKPNGCRKWETTFKKTSD DGEVYYSEEAKKKVEVLDTDYKSYAVIYATRVKDGRTLHMMRLYSRSPVSPAATAIFRKLGERNYTDEMVA MLPRQEECTVDEV

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 85% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/ μ g of the protein as determined by the LAL method.
Formulation:	If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μ g/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Siderocalin-like lipocalin tightly binding a variety of bacterial ferric siderophores, also binds long-chain unsaturated fatty acids such as linoleic acid, oleic acid, arachidonic acid and, with a lower affinity, long chain

A DRUG SCREENING EXPERT

saturated fatty acids such as stearic acid. May act as an antibacterial factor, through dual ligand specificity, both as a siderophore-sequestering molecule and a lysophosphatidic acid (LPA) sensor.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481