

GSTP1 Protein, Cricetus longicaudatus, Recombinant (His)

General Information

Synonyms:	GSTP1; Glutathione S-transferase P; GST class-pi
Protein Construction:	1-210 aa
Species:	Chinese hamster
Expression Host:	P. pastoris (Yeast)
Accession:	P46424
Molecular Weight:	26.3 kDa (predicted)
AA Sequence:	MPPYTIVYFPVGRCEAMRILLADQGQSWKEEVVTVETWRKGLKSTCLYGQLPKFEDGDLTYQSNAILRHLGRSLGLYGKDKQREAAALVDMVNDGVEDLRCKYITLIYTKYEEGKDDYVKALPGHLKPFETLLSQNQGGKAFIVGDQISFVDYNLLDLLLIHQVLAPGCLDNFPLLSAYVARLSARPKIKAFLLSSPDHVNRPINGNGKQ

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 85% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles. Involved in the formation of glutathione conjugates of both prostaglandin A2 (PGA2) and prostaglandin J2 (PGJ2).

A DRUG SCREENING EXPERT

Participates in the formation of novel hepoxilin regioisomers. Regulates negatively CDK5 activity via p25/p35 translocation to prevent neurodegeneration.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481