

## PLA2G15 Protein, Canine, Recombinant (His & Myc)

### General Information

Synonyms:	1-O-acylceramide synthase (ACS);LYPLA3;Lysosomal phospholipase A and acyltransferase; PLA2G15;Phospholipase A2 group XV;LCAT-like lysophospholipase (LLPL);Lysophospholipase 3;Lysosomal phospholipase A2 (LPLA2)
Protein Construction:	32-408 aa
Species:	Canine
Expression Host:	Baculovirus Insect Cells
Accession:	Q6XPZ3
Molecular Weight:	46.8 kDa (predicted)
AA Sequence:	RRPPVVLVPGDLGNQLEAKLDKPTVVHYLCSKRTESYFTLWLNLELLLPIIDCWIDNIRLVYNRTSRATQFPDG VDVRVPGFGKTFSLFLDPSKSSVGSYFHTMVESLVDWGWYIRGEDVRGAPYDWRRAPNENGPYFLALREMI EMYQLYGGPVVLAHSMGNMYTLYFLQRQPQAWKNKYIQAFVALGAPWGGVAKTLRVLASGDNNRIPVIR PLKIREQQRSAVSTSWLLPYNYTWSPEKIFVHTPTANYTLRDYHQFFQDIGFKDGWLMRQDTEGLVEAMVPP GVPLHCLYGTGVPTPDSFYYESFPDRDPKICFGDGDGTVNLQSALQCQAWRGHQEHQVSLQALPGSEHIEM LANATTLAYLKRVLG

### QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 85% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol.If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

### Preparation and Storage

#### Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

#### Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

#### Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

Has dual calcium-independent phospholipase and O-acyltransferase activities with a potential role in glycerophospholipid homeostasis and remodeling of acyl groups of lipophilic alcohols present in acidic cellular compartments. Catalyzes hydrolysis of the ester bond of the fatty acyl group attached at sn-1 or sn-2 position of phospholipids (phospholipase A1 or A2 activity) and transfer it to the hydroxyl group at the first carbon of lipophilic alcohols (O-acyltransferase activity). Among preferred fatty acyl donors are phosphatidylcholines, phosphatidylethanolamines, phosphatidylglycerols and phosphatidylserines. Favors sn-2 over sn-1 deacylation of unsaturated fatty acyl groups of phosphatidylcholines and phosphatidylethanolamines. Among preferred fatty acyl acceptors are natural lipophilic alcohols including short-chain ceramide N-acetyl-sphingosine (C2 ceramide), alkylacylglycerols, monoacylglycerols, and acylethanolamides such as anandamide and oleoylethanolamide. Selectively hydrolyzes the sn-1 fatty acyl group of truncated oxidized phospholipids and may play a role in detoxification of reactive oxidized phospholipids during oxidative stress. Required for normal phospholipid degradation in alveolar macrophages with potential implications in pulmonary surfactant clearance. At neutral pH, hydrolyzes the sn-1 fatty acyl group of the lysophosphatidylcholines.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481