

Cutinase Protein, Fusarium solani subsp. Cucurbitae, Recombinant

General Information

| | |
|-----------------------|---|
| Synonyms: | Cutinase;Cutin hydrolase;CUTA |
| Protein Construction: | 17-230 aa |
| Species: | Neocosmosporum cucurbitae |
| Expression Host: | E. coli |
| Accession: | Q99174 |
| Molecular Weight: | 22.4 kDa (predicted) |
| AA Sequence: | LPTSNPAQEARQLGRTRDDLINGNSASCADVIFIYARGSTETGNLGLGPSIASNLESFAFGTDGVWIIQGVG GAYRATLGDNALPRGTSSAAIREMLGLFQQANTKCPDATLIAGGYSQGAALAAASIEDLDSAIRDKIAGTVLF GYTKNLQNRGRIPNYPADRTKVFVCGDLVCTGSLIVAAPHLAYGPDARGPAPEFLIEKVRVAVRGSAA |

QC Testing

| | |
|----------------------|--|
| Biological Activity: | Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first. |
| Purity: | > 90% as determined by SDS-PAGE. |
| Endotoxin: | < 1.0 EU/μg of the protein as determined by the LAL method. |
| Formulation: | If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0. |

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Catalyzes the hydrolysis of cutin, a polyester that forms the structure of plant cuticle. Allows pathogenic fungi to penetrate through the cuticular barrier into the host plant during the initial stage of the fungal infection.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481