

HTR1F Protein, Human, Recombinant (His & SUMO)

General Information

Synonyms:	HTR1F;5-HT-1F;5-HT1F;HTR1EL;Serotonin receptor 1F;5-hydroxytryptamine receptor 1F
Protein Construction:	1-366 aa
Species:	Human
Expression Host:	E. coli
Accession:	P30939
Molecular Weight:	60.2 kDa (predicted)
AA Sequence:	MDFLNSSDQNLTSSELLNRMPKILVSLTSLGLALMTTINSLVIAAIIIVTRKLHHPANYLICSLAVTDFLVAVLV MPFSIVYIVRESWIMGQVVCDIWLSVDITCCTCSILHLSAIALDRYRAITDAVEYARKRTPKHAGIMITIVWIISVFI SMPPLFWRHQGTSRDDECIKHDHIVSTIYSTFGAFYIPLALILILYKIIYRAAKTLYHKRQASRIAKEEVNGQVLL ESGEKSTKSVSTSYVLEKSLSDPSTDFDKIHSTVRSRSEFKHEKSWRRQKISGTRERKAATTLGLILGAFVICWL PFFVKELVVNVCCKCKISEEMSNFLAWLGYLNSLINPLIYTIFNEDFKKAFQKLVRCRC

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 85% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from Tris/PBS-based buffer, 6% Trehalose, pH 8.0

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

G-protein coupled receptor for 5-hydroxytryptamine (serotonin). Also functions as a receptor for various alkaloids and psychoactive substances. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors, such as adenylate

cyclase. Signaling inhibits adenylate cyclase activity.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481