

## Human adenovirus C serotype 2 Early E3 18.5 kDa glycoprotein (His & Myc)

### General Information

Synonyms:	E3-19K;E3gp 19 kDa (E19);Early E3 18.5 kDa glycoprotein;GP19K
Protein Construction:	18-123 aa
Species:	HAdV-2
Expression Host:	E. coli
Accession:	P68978
Molecular Weight:	20.0 kDa (predicted)
AA Sequence:	AKKVEFKEPACNVTFKSEANECTTLIKCTTEHEKLIIRHKDKIGKYAVYAIWQPGDTNDYNVTVFQGENRKTFM YKFPFYEMCDITMYMSKQYKLWPPQKLENTG

### QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

### Preparation and Storage

#### Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

#### Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

#### Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

Binds and retains class I heavy chains in the endoplasmic reticulum during the early period of virus infection, thereby impairing their transport to the cell surface. Also delays the expression of class I alleles that it cannot affect by direct retention. Binds transporters associated with antigen processing (TAP) and acts as a tapasin

## A DRUG SCREENING EXPERT

---

inhibitor, preventing class I/TAP association. In consequence, infected cells are masked for immune recognition by cytotoxic T-lymphocytes.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481