

MGAT4A Protein, Human, Recombinant (His & Myc)

General Information

Synonyms:	3-mannosyl-glycoprotein 4-beta-N-acetylglucosaminyltransferase A;N-glycosyl-oligosaccharide-glycoprotein N-acetylglucosaminyltransferase IVa (GlcNAc-T IVa;GnT-IVa;N-acetylglucosaminyltransferase IVa);MGAT4A;UDP-N-acetylglucosamine: alpha-1,3-D-mannoside beta-1,4-N-acetylglucosaminyltransferase IVa;Alpha-1,3-mannosyl-glycoprotein 4-beta-N-acetylglucosaminyltransferase A
Protein Construction:	93-535 aa
Species:	Human
Expression Host:	HEK293 Cells
Accession:	Q9UM21
Molecular Weight:	56 kDa (predicted)
AA Sequence:	LLKELTSKSLQVPSIYYHLPHLLKNEGSLQPAVQIGNGRGTGVSIVMGIPTVKREVKSYLIETLHSLIDNLYPEEKLD DCVIVVFIGETDIDYVHGVVANLEKEFSKEISSGLVEISPPESYYPDLTNLKETFGDSKERVRWRWKQNLDYCFL MMYAQEKGIYYIQLEDDIIVKQNYFNTIKNFALQLSSEEWMILEFSQLGFIGKMFQAPDLTLIVEFIFMFYKEKPI DWLLDHILWVKVCNPEKDAKHCDRQKANLRIRFRPSLFQHVGLHSSLGKIQKLTDKDYMKPLLLKIHVNP PPAEVSTSLKVYQGHTEKTYMGEDFFWAITPIAGDYILFKFDKPVNVESYLFHSGNQEHPGDILLNTTVEVLPFKSE GLEISKETKDKRLEDGYFRIGKFENGVAEGMVDPSLNPISAFRLSVIQNSAVWAILNEIHIKKATN

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 85% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months.

Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Glycosyltransferase that participates in the transfer of N-acetylglucosamine (GlcNAc) to the core mannose residues of N-linked glycans. Catalyzes the formation of the GlcNAc β 1-4 branch on the GlcNAc β 1-2Man α 1-3 arm of the core structure of N-linked glycans. Essential for the production of tri- and tetra-antennary N-linked sugar chains. Involved in glucose transport by mediating SLC2A2/GLUT2 glycosylation, thereby controlling cell-surface expression of SLC2A2 in pancreatic beta cells.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481