

Aquaporin-5/AQP5 Protein, Human, Recombinant (His)

General Information

Synonyms: AQP-5;Aquaporin-5;AQP5

Protein Construction: 1-265 aa

Species: Human

Expression Host: E. coli

Accession: P55064

Molecular Weight: 31.1 kDa (predicted)

AA Sequence:

MKKEVCSVAFLKAVFAEFLATLIFVFFGLGSALKWPSALPTILQIALAFGLAIGTLAQAALGPVSGGHINPAITLAL
LVGNQISLLRAFFYVAAQLVGAIAGAGILYGVAPLNARGNLAVNALNNNTTQGGQAMVVELILTFQLALCIFAST
DSRRTSPVGPALSIGLSVTLGHLVGIYFTGCSMNPARSFGPAVVMNRFSPAHWVFWVGPIVGAVLAAILYFY
LLFPNSLSLSEVAIIKGTYPDEDWEEQREERKKTMELTTR

QC Testing

Biological Activity: Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.

Purity: > 85% as determined by SDS-PAGE.

Endotoxin: < 1.0 EU/μg of the protein as determined by the LAL method.

Formulation: If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Forms a water-specific channel. Plays an important role in fluid secretion in salivary glands. Required for TRPV4

A DRUG SCREENING EXPERT

activation by hypotonicity. Together with TRPV4, controls regulatory volume decrease in salivary epithelial cells. Seems to play a redundant role in water transport in the eye, lung and in sweat glands.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481