

MAP2K3 Protein, Human, Recombinant (His & SUMO)

General Information

Synonyms:	PRKMK3;MAPKK 3;SKK2;MKK3;Dual specificity mitogen-activated protein kinase kinase 3;MAP kinase kinase 3;Stress-activated protein kinase kinase 2 (SAPK kinase 2;SAPKK-2;SAPKK2);MEK3;MAP2K3;MAPK/ERK kinase 3 (MEK 3)
Protein Construction:	1-347 aa
Species:	Human
Expression Host:	E. coli
Accession:	P46734
Molecular Weight:	55.3 kDa (predicted)
AA Sequence:	MESPASSQPASMPQSKGKSKRKKDLRISCMSPAPNPPTPPRNLDSTRFITIGDRNFEVEADDLVTISELGRG AYGVVEKVRHAQSGTIMAVKRIRATVNSQEQRLLMDLDINMRTVDCFYTVTFYGALFREGDVVICMELMDT SLDKFYRKVLDKNMTIPEDILGEIAVSIVRALEHLHSLKSLVIHRDVKPSNVLINKEGHVKMCDFGISGYLVDSVA KTMDAGCKPYMAPERINPELNQKGYNVKSDVWSLGITMIEMAILRFPYESWGTGPFQQLKQVVEEPPQLPAD RFSPEFVDFTAQCLRKNPAERMSYLELMEHPFFTLHKTKKTDIAAFVKEILGEDS

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Tris-based buffer, 50% glycerol

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Dual specificity kinase. Is activated by cytokines and environmental stress in vivo. Catalyzes the concomitant

phosphorylation of a threonine and a tyrosine residue in the MAP kinase p38. Part of a signaling cascade that begins with the activation of the adrenergic receptor ADRA1B and leads to the activation of MAPK14.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481