

GABRP Protein, Human, Recombinant (His)

General Information

Synonyms:	GABRP; Gamma-aminobutyric acid receptor subunit pi; GABA(A) receptor subunit pi (GABAAR subunit pi)
Protein Construction:	17-242 aa
Species:	Human
Expression Host:	E. coli
Accession:	O00591
Molecular Weight:	28.9 kDa (predicted)
AA Sequence:	ERMCIQGSQFNVEVGRSDKLSLPGFENLTAGYNKFLRPNFGGPEVQIALTLDIASISSISESNMDYTATIYLRQR WMDQRLVFEGNKSFTLDARLVEFLWVPDTYIVESKKSFLHEVTVGNRLIRLFSNGTVLYALRITTTVACNMDLS KYPMDTQTCKLQLESWGYDGDNDVEFTWLRGNDVSRGLEHLRLAQYTIERYFTLVTRSQQETGNYTRLVLQFE LRRNV

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 85% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

A DRUG SCREENING EXPERT

GABA, the major inhibitory neurotransmitter in the vertebrate brain, mediates neuronal inhibition by binding to the GABA/benzodiazepine receptor and opening an integral chloride channel. In the uterus, the function of the receptor appears to be related to tissue contractility. The binding of this $\alpha 1$ subunit with other GABA(A) receptor subunits alters the sensitivity of recombinant receptors to modulatory agents such as pregnanolone.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481