

Gasdermin-C Protein, Human, Recombinant (His & Myc)

General Information

Synonyms:	Gasdermin-C;MLZE;GSDMC;Melanoma-derived leucine zipper-containing extranuclear factor
Protein Construction:	1-508 aa
Species:	Human
Expression Host:	Baculovirus Insect Cells
Accession:	Q9BYG8
Molecular Weight:	61.7 kDa (predicted)
AA Sequence:	MPSMLERISKNLVKEIGSKDLTPVKYLLSATKLRQFVILRKKKDSRSSFWEQSDYVPVEFSLNDILEPSSSVLETV VTGPFHFSDDIMIQQHKADMGVNVGIEVSVSGEASVDHGCSLEFQIVTIPSPNLEDFQKRKLLDPEPSFLKECRR RGDNLYVVTEAVELINNTVLYDSSSVNIGKIALWITYGKGQGGESLRVKKKALTQKGMVMAYKRKQLVIK EKAILISDDDEQRTFQDEYEISEMVGYCAARSEGLLPSFHTISPTLFNASSNDMKLKPFLTQQFLSGHLPKYE QVHILPVGRIEPPFWQNFKHLQEEVFQKIKTLAQLSKDQDVMFYSLAMLRDRGALQDLMNMLELDSSGHL DGPGGAILKKLQQDSNHAWFNPDPILYLLEAIMVLSDFQHDLLACSMKRILLQQQELVRSILEPNFRYPWSI PFTLKPPELLAPLQSEGLAITYGLLEECGLRMELDNPRSTWDVEAKMPLSALYGTLSLLQLAEA

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 85% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Tris-based buffer, 50% glycerol

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

A DRUG SCREENING EXPERT

This form constitutes the precursor of the pore-forming protein: upon cleavage, the released N-terminal moiety (Gasdermin-C, N-terminal) binds to membranes and forms pores, triggering cell death.; Pore-forming protein that causes membrane permeabilization and pyroptosis. Released upon cleavage and binds to membrane inner leaflet lipids. Homooligomerizes within the membrane and forms pores of 10-15 nanometers (nm) of inner diameter, triggering pyroptosis. The functional mechanisms and physiological proteases that cleave and activate this pore-forming protein are unknown (Probable).

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481