

IL12RB2 Protein, Human, Recombinant (His)

General Information

Synonyms:	IL-12R subunit beta-2;IL-12RB2;IL-12R-beta-2;Interleukin-12 receptor subunit beta-2;IL12RB2;IL-12 receptor subunit beta-2
Protein Construction:	24-622 aa
Species:	Human
Expression Host:	P. pastoris (Yeast)
Accession:	Q99665
Molecular Weight:	69.7 kDa (predicted)
AA Sequence:	KIDACKRGDVTVKPSHVILLGSTVNITCSLKPRQGCFFHYSRRNKLILYKFDRRINFHHGHSLSNSQVTGLPLGTTL FVCKLACINSDEIQICGAEIFVGVAPQPNLSCIQKGEQGTVACTWERGRDTHLYTEYTLQLSGPKNLTWQK QCKDIYCDYLDLFGINLTPESPESNFTAKVTAVNSLGSSSLPSTFTFLDIVRPLPPWDIRIKFQKASVSRCTLYWR DEGLVLLNRLRYPNSRLWNMNVNVTAKGRHLLDLKPFTEYEFQISSKLHLYKGSWSDWSESLRAQTPEE EPTGMLDVWYMKRHIDYSRQQLSLFWKNLSVSEARGKILHYQVTLQELTGGKAMTQNTGHTSWTTVIPRTG NWAVAVSAANSKGSLLPTRINIMNLCEAGLLAPRQVSANSEGMNLTWQPPRKDPSAVQEYVVEWRELH PGGDTQVPLNWLRSRPNVSAISENIKSYICYEIRVYALSGDQGGCSSILGNSKHKAPLSGPHINAITEEKGSIL ISWNSIPVQEQMGCLLHYRIYWKERDSNSQPQLCEIPYRVSQNSHPINSLQPRVTYVLWMTALTAAGESSHG NEREFCLQGKAN

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Tris-based buffer, 50% glycerol

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Receptor for interleukin-12. This subunit is the signaling component coupling to the JAK2/STAT4 pathway. Promotes the proliferation of T-cells as well as NK cells. Induces the promotion of T-cells towards the Th1 phenotype by strongly enhancing IFN-gamma production.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481