

## GBA Protein, Human, Recombinant

## General Information

Synonyms:	Cholesterol glucosyltransferase (SGTase); Glucosylceramidase beta 1; Lysosomal acid glucosylceramidase; Lysosomal glycosylceramidase; Beta-glucosylceramidase 1; Lysosomal acid GCase; GBA1; Imiglucerase; Beta-glucocerebrosidase (Beta-GC); GC; D-glucosyl-N-acylsphingosine glucohydrolase; Lysosomal cholesterol glucosyltransferase; GBA; Lysosomal galactosylceramidase; Acid beta-glucosidase; GLUC; Alglucerase; Cholesteryl-beta-glucosidase
Protein Construction:	40-536 aa
Species:	Human
Expression Host:	E. coli
Accession:	P04062
Molecular Weight:	55.6 kDa (predicted)
AA Sequence:	ARPCIPKSFYSSVVCVFNATYCDSDPPTFPALGTFSTRYESTRSGRRMELSMGPIQANHTGTGLLLTLQPEQK FQKVKGFGGAMTDAAALNILALSPPAQNLLLSYFSEEIGYNIIRVPMASCFDSIRTYTYADTPDDFQLHNFS LPEEDTKLKIPLIHRALQLAQRPVSLASPWTSPWLKTNGAVNGKGLKGPQGDYHQTWARYFVKFLDAYA EHKLQFWAVTAENEPSAGLLSGYPFQCLGFTPEHQRFIARDLGPTLANSTHHNVRLMLDDQRLLLPHWAK VVLTDPEAAKYVHGIADVHWYLDLFLAPAKATLGETHRLFPNTMLFASEACVGSKFWEQSVRLGSDWRGMQYS HSIITSLLYHVVGWTDWNLALNPEGGPNWVRNFVDSPIIVDITKDTFYKQPMFYHLGHFSKFIPEGSQRVGLV ASQKNDLDAVALMHPDGSVVVVLNRSSKDVPLTIKDPAVGFLETISPYSIHTYLWRRQ

## QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 85% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Tris-based buffer, 50% glycerol

## Preparation and Storage

## Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

## Stability &amp; Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

*Actual storage temperature shall be subject to the COA.*

## Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

Glucosylceramidase that catalyzes, within the lysosomal compartment, the hydrolysis of glucosylceramide/GlcCer into free ceramide and glucose. Thereby, plays a central role in the degradation of complex lipids and the turnover of cellular membranes. Through the production of ceramides, participates in the PKC-activated salvage pathway of ceramide formation. Also plays a role in cholesterol metabolism. May either catalyze the glucosylation of cholesterol, through a transglucosylation reaction that transfers glucose from glucosylceramide to cholesterol. The short chain saturated C8:0-GlcCer and the mono-unsaturated C18:0-GlcCer being the most effective glucose donors for that transglucosylation reaction. Under specific conditions, may alternatively catalyze the reverse reaction, transferring glucose from cholesteryl-beta-D-glucoside to ceramide. Finally, may also hydrolyze cholesteryl-beta-D-glucoside to produce D-glucose and cholesterol.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481