

## MRGPRX2 Protein, Human, Recombinant (His)

### General Information

Synonyms: MRGX2;Mas-related G-protein coupled receptor member X2;MRGPRX2

Protein Construction: 1-330 aa

Species: Human

Expression Host: In vitro E. coli

Accession: Q96LB1

Molecular Weight: 43.1 kDa (predicted)

AA Sequence:

MDPTTPAWGTESTTVNGNDQALLLFCGKETLIPVFLILFIALVGLVGNFVLWLLGFRMRRNAFSVYVLSLAGA  
DFLFLCFQIINCLVYLSNFFCSISINFPSTTVMTCAYLAGLSMLSTVSTERCLSVLWPIWYRCRRPRHLSAVVC  
VLLWALSLLLSILEGKFCGFLFSDGDSGWCQTFDFITAAWLIFLMVLCGSSLALLVRILCGSRGLPLTRLYLTILL  
TVLVFLLCGLPFGIQWFLILWIWKDSDVLFCHIHPVSVVLSLSSANPIIYFFVGSFRKQWRLQQPILKLALQR  
ALQDIAEVDHSEGCFRQGTPEMSRSSLV

### QC Testing

Biological Activity: Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.

Purity: > 90% as determined by SDS-PAGE.

Endotoxin: < 1.0 EU/μg of the protein as determined by the LAL method.

Formulation: If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

### Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

## A DRUG SCREENING EXPERT

---

Mast cell-specific receptor for basic secretagogues, i.e. cationic amphiphilic drugs, as well as endo- or exogenous peptides, consisting of a basic head group and a hydrophobic core. Recognizes and binds small molecules containing a cyclized tetrahydroisoquinoline (THIQ), such as non-steroidal neuromuscular blocking drugs (NMBDs), including tubocurarine and atracurium. In response to these compounds, mediates pseudo-allergic reactions characterized by histamine release, inflammation and airway contraction. Acts as a receptor for a number of other ligands, including peptides and alkaloids, such as cortistatin-14, proadrenomedullin N-terminal peptides PAMP-12 and, at lower extent, PAMP-20, antibacterial protein LL-37, PMX-53 peptide, beta-defensins, and complanadine A.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481