

MAGEA3 Protein, Human, Recombinant (His & Myc)

General Information

Synonyms:	Antigen MZ2-D;Cancer/testis antigen 1.3 (CT1.3);MAGE3;MAGE-3 antigen;MAGEA3;Melanoma-associated antigen 3
Protein Construction:	1-314 aa
Species:	Human
Expression Host:	E. coli
Accession:	P43357
Molecular Weight:	39.8 kDa (predicted)
AA Sequence:	MPLEQRSQHCKPEEGLEARGEALGLVGAQAPATEEQEAASSSSTLVEVTLGEVPAAESPDPQPQSPQGASSLP TTMNYPLWSQSYEDSSNQEEEGPSTFPDLESEFQAALSRKVAELVHFLLLKYRAREPVTKAEMLGSSVGNWQ YFFPVIFSKASSLQLVFGIELMEVDPIGHLYIFATCLGLSYDGLLDGNQIMPKAGLLIIVLAIAREGDCAPEEKIW EELSVLEVFEGREDSILGDPKLLTQHFVQENYLEYRQVPGSDPACYEFLWGPRLVETSIVKVLHMHMVKISG GPHISYPPLHEWVLRGEE

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 85% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Tris-based buffer, 50% glycerol

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Activator of ubiquitin ligase activity of RING-type zinc finger-containing E3 ubiquitin-protein ligases that acts as a repressor of autophagy. May enhance ubiquitin ligase activity of TRIM28 and stimulate p53/TP53 ubiquitination

A DRUG SCREENING EXPERT

by TRIM28. Proposed to act through recruitment and/or stabilization of the Ubl-conjugating enzyme (E2) at the E3: substrate complex. May play a role in embryonal development and tumor transformation or aspects of tumor progression. In vitro promotes cell viability in melanoma cell lines. Antigen recognized on a melanoma by autologous cytolytic T-lymphocytes.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481