

GPR107 Protein, Human, Recombinant (His & Myc)

General Information

Synonyms:	GPR107;KIAA1624;Protein GPR107;Lung seven transmembrane receptor 1;LUSTER1
Protein Construction:	40-263 aa
Species:	Human
Expression Host:	E. coli
Accession:	Q5VW38
Molecular Weight:	32.4 kDa (predicted)
AA Sequence:	RVHHLALKDDVRHKVHLNTEFGFFKGYMNVNVSLSLNEPEDKDVTIGFSLDRTKNDGFSSYLDEDVNYCILK KQSVSVTLILDISRSEVRVKSPPEAGTQLPKIIFSRDEKVLGQSQEPNPNPASAGNQTQKTQDGGKSKRSTVD SKAMGEKSFVHNNGGAVSFQFFFNISTDDQEGLYSLYFHKCLGKELPSDKFTFSLDIEITEKNPDSYLSAGEIP LP

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

Preparation and Storage

Reconstitution:	Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.
Stability & Storage:	Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots. <small>Actual storage temperature shall be subject to the COA.</small>
Shipping:	In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Has been proposed to act as a receptor for neuronostatin, a peptide derived from the somatostatin/SST precursor.

A DRUG SCREENING EXPERT

Involved in blood sugar regulation through the induction of glucagon in response to low glucose.; (Microbial infection) Required for intoxication by *Pseudomonas aeruginosa* exotoxin A and *Campylobacter jejuni* CDT. May contribute to the retrograde transport of bacterial toxins, including cholera toxin, from the trans-Golgi network to the endoplasmic reticulum.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481