

Sialidase-3 Protein, Human, Recombinant (His)

General Information

Synonyms:	N-acetyl-alpha-neuraminidase 3;Sialidase-3;Membrane sialidase;Ganglioside sialidasedis; NEU3
Protein Construction:	2-428 aa
Species:	Human
Expression Host:	E. coli
Accession:	Q9UQ49
Molecular Weight:	52.1 kDa (predicted)
AA Sequence:	EEVTTCSFNSPLFRQEDDRGITYRIPALLYIPPTHTFLAFAEKRSTRRDEDALHLVLRRLRIGQLVQWGPLKPL MEATLPGHRTMNPVWEQKSGCVLFFICVRGHVTERQQIVSGRNAARLCFIYSQDAGCSWSEVRDLTEEVI GSELKHWATFAVGPVGHGIQLQSGRLVIPAYTYIIPSWFFCFQLPCKTRPHSLMIYSDDLGVTWHHGRLIRPMV TVECEVAEVTGRAGHPVLYCSARTPNRCRAEALSTDHGEGFQRLALSRLCEPPHGCQGSVVSFRPLEIPHRC QDSSSKDAPTIQOSSPGSSLRLEEEAGTPSESWLLYSHPTSRKQRVDLGIYLNQTPLEAACWSRPWILHCGPC GYSDLAALIEEGLFGCLFECGKQECEQIAFRLFTHREILSHLQGDCTSPGRNPSQFKSN

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Tris-based buffer, 50% glycerol

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Exo-alpha-sialidase that catalyzes the hydrolytic cleavage of the terminal sialic acid (N-acetylneuraminic acid,

A DRUG SCREENING EXPERT

Neu5Ac) of a glycan moiety in the catabolism of glycolipids, glycoproteins and oligosaccharides. Displays high catalytic efficiency for gangliosides including alpha-(2->3)-sialylated GD1a and GM3 and alpha-(2->8)-sialylated GD3. Plays a role in the regulation of transmembrane signaling through the modulation of ganglioside content of the lipid bilayer and by direct interaction with signaling receptors, such as EGFR. Desialylates EGFR and activates downstream signaling in proliferating cells. Contributes to clathrin-mediated endocytosis by regulating sorting of endocytosed receptors to early and recycling endosomes.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481