

TLR7 Protein, Human, Recombinant (His & Myc)

General Information

Synonyms:	Toll-like receptor 7;TLR7
Protein Construction:	861-1049 aa
Species:	Human
Expression Host:	E. coli
Accession:	Q9NYK1
Molecular Weight:	29.5 kDa (predicted)
AA Sequence:	HLYFWDVWYIYHFCKAKIKGYQRLISPDCCYDAFIVYDTKDAVTEWVLAELVAKLEDPREKHFNLCEERDW LPGQPVLENLSQSIQLSKKTVMFMTDKYAKTENFKIAFYLSHQRLMDEKVDVILIFLEKPFQKSKFLQLRKRLCG SSVLEWPTNPQAHFYFWQCLKNALATDNHVAYSQVFKETV

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 85% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Tris-based buffer, 50% glycerol

Preparation and Storage

Reconstitution:
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Endosomal receptor that plays a key role in innate and adaptive immunity. Controls host immune response against pathogens through recognition of uridine-containing single strand RNAs (ssRNAs) of viral origin or guanosine analogs. Upon binding to agonists, undergoes dimerization that brings TIR domains from the two molecules into direct contact, leading to the recruitment of TIR-containing downstream adapter MYD88 through homotypic interaction. In turn, the Myddosome signaling complex is formed involving IRAK4, IRAK1, TRAF6, TRAF3

A DRUG SCREENING EXPERT

leading to activation of downstream transcription factors NF-kappa-B and IRF7 to induce proinflammatory cytokines and interferons, respectively.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481