

pylS Protein, Methanosarcina barkeri, Recombinant (His)

General Information

Synonyms:	pylS;Pyrrolysyl-tRNA synthetase (PylRS);Pyrrolysine--tRNA ligase;Pyrrolysine--tRNA(Pyl) ligase
Protein Construction:	1-419 aa
Species:	Methanosarcina barkeri
Expression Host:	E. coli
Accession:	Q46E77
Molecular Weight:	51.8 kDa (predicted)
AA Sequence:	MDKKPLDVLISATGLWMSRTGTLHKIKHYEVSRSKIYIEMACGDHLVVNNSRSCRTARAFRHHKYRKTCKRCR VSDDEDINNFLTRSTEGKTSVKVKVVSAPKVKKAMPKSVSRAPKPLENPVSAKASTDTSRSPSPAKSTPNSPV PTSAPAPSLTRSQLDRVEALLSPEDKISLNIAPPFRELESELVTRRKNDFQRLYTNDREDYLGKLERDITKFFVDR DFLEIKSPILIPAEYVERMGINNDTELSKQIFRVDKNLCLRPMLAPTLNYLRKLDRLIPDPIKIFEVGPCYRKESD GKEHLEFTMVNFQCMGSGCTRENLES LIKEFLDYLEIDFEIVGDSCMVYGDTLDIMHGDLESSAVVGPVPLD REWGIDKPWIGAGFGLERLLKVMHGFKNIKRASRSSESYNGISTNL

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 85% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Catalyzes the attachment of pyrrolysine to tRNA(Pyl). Pyrrolysine is a lysine derivative encoded by the termination codon UAG.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481