

## AKR1C3 Protein, Mouse, Recombinant (His & Myc)

### General Information

Synonyms: Aldo-keto reductase family 1 member C13;Akr1c13

Protein Construction: 1-323 aa

Species: Mouse

Expression Host: E. coli

Accession: Q8VC28

Molecular Weight: 44.5 kDa (predicted)

AA Sequence: MSSKQHCVKLNDGHLIPALGFGTYKPKEVPKSKSLEAACLALDVGYRHVDTAYAYQVEEEIGQAIQSKIKAGV  
VKREDLFITTKLWCTCFRPELVKPALEKSLKQLDLYVDLYIMHYPVPMKSGDNDFPVNEQGSLLDVTDFCD  
TWERLEECKDAGLVKISIGVSNFNHRQLERILNKPGLKYKPCNQQVECHLYLNQRKLLDYCESKDIVLVAYGAL  
GTQRYKEWVDQNSPVLNLPVLCDAKKNKRSPALIALRYLIQRGIVPLAQSFKENEMRENLQVFGFQLSPE  
DMKTLDGLNKNFRYLPAEFLVDHPEYPFVEEY

### QC Testing

Biological Activity: Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.

Purity: > 85% as determined by SDS-PAGE.

Endotoxin: < 1.0 EU/μg of the protein as determined by the LAL method.

Formulation: If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

### Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

## A DRUG SCREENING EXPERT

---

Catalyzes the dehydrogenation of 17-beta-hydroxysteroids. May also exhibit significant activity with a variety of cyclic and alicyclic alcohols. Uses both NAD and NADP, but the activity is much greater with NAD than with NADP.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481