

BAG3 Protein, Mouse, Recombinant (His)

General Information

Synonyms:	BAG family molecular chaperone regulator 3; Bag3; Bcl-2-associated athanogene 3; BAG-3; Bis; Bcl-2-binding protein Bis
Protein Construction:	2-577 aa
Species:	Mouse
Expression Host:	P. pastoris (Yeast)
Accession:	Q9JLV1
Molecular Weight:	63.7 kDa (predicted)
AA Sequence:	SAATQSPMMQMASGNGASDRDPLPPGWEIKIDPQTGWPFVDHNSRTTWTNDPRVPPEGPKDTASSANG PSRDGSRLLPIREGHPYIYQLRPGYIPIVLEHGSERQPHLFHAYSQPGVQRFRTAAAATPQRSQSPLRGG MTEAAQTDKQCGQMPATATATAAQPPTAHGPERSQSPAASDCSSSSSSASLPSSGRSSLGSHQLPRGYIPI VIHEQNITRPAAQPSFHQAQKTHYPAQQGEYQPQQPVYHKIQGDDWEPRPLRAASPFVSPVREGASSREGSP ARSGTPVHCPSPVHTVDRPQPMTHREPPVTPENKPKESKPGPAGPDLPPGHIPIQVIRREADSKPVSQK SPPPAEKVEVKVSSAPIPCPSPSPAPSAVPSPPKNVAAEQKAAPSPAPAEPAPKSGEAEPPKHPGVLKVEA ILEKVQGLEQAVDSFEGKTKDKKYLMIIEYLTKELLALDSVDEPGRADVRQARRDGVKRVQVILEKLEQKVIDV PGVQVYELQPSNLEAEQPLQEIMGAVVADKDKKGPENKDPQTESQLEAKAATPPNPSNPADSAGNLVA P

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Tris-based buffer, 50% glycerol

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months.

Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Co-chaperone for HSP70 and HSC70 chaperone proteins. Acts as a nucleotide-exchange factor (NEF) promoting the release of ADP from the HSP70 and HSC70 proteins thereby triggering client/substrate protein release. Nucleotide release is mediated via its binding to the nucleotide-binding domain (NBD) of HSPA8/HSC70 where as the substrate release is mediated via its binding to the substrate-binding domain (SBD) of HSPA8/HSC70. Has anti-apoptotic activity. Plays a role in the HSF1 nucleocytoplasmic transport.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481