

CCL4 Protein, Mouse, Recombinant (His)

General Information

Synonyms:	Small-inducible cytokine A4;SIS-gamma;C-C motif chemokine 4;Ccl4;Immune activation protein 2 (ACT-2;ACT2);Scya4;Mip1b;Macrophage inflammatory protein 1-beta (MIP-1-beta); Protein H400
Protein Construction:	24-92 aa
Species:	Mouse
Expression Host:	E. coli
Accession:	P14097
Molecular Weight:	11.8 kDa (predicted)
AA Sequence:	APMGSDPPTSCCFSYTSRQLHRFSVMDYYETSSLCSKPAVVFLTKRGRQICANPSEPWVTEYMSDLELN

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Tris-based buffer, 50% glycerol

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Monokine with inflammatory and chemokinetic properties.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481