

HSPA5 Protein, Mouse, Recombinant (His)

General Information

Synonyms:	78 kDa glucose-regulated protein (GRP-78);Heat shock protein family A member 5;Heat shock protein 70 family protein 5 (HSP70 family protein 5);Endoplasmic reticulum chaperone BiP;Binding-immunoglobulin protein (BiP);Hspa5;Grp78;Immunoglobulin heavy chain-binding protein
Protein Construction:	20-655 aa
Species:	Mouse
Expression Host:	P. pastoris (Yeast)
Accession:	P20029
Molecular Weight:	72.5 kDa (predicted)
AA Sequence:	EEEDKKEDVGTVVGIDLGTTYSCVGVFKNGRVEIANDQGNRITPSYVAFTPEGERLIGDAAKNQLTSNPENTV FDAQRLIGRTWNDPSVQQDIKFLPFKVVVEKTKPKYIQVDIGGGQTKTFAPEEISAMVLTKMKETAAYLGKKVT HAVVTVPAYFNDAQRQATKDAGTIAGLNVMRIINEPTAAAIAYGLDKREGEKNILVFDLGGGTFDVSLLTIDNG VFEVVATNGDTHLGGEDFDQRMVMEHFILYKKTGKDV RKDNRAVQKLRREVEKAKRALSSQHARIEIESFF EGEDFSETLTRAKEELNMDLFRSTMKPVQKVLSDSKDIDEIVLGGSTRIPKIQQLVKEFFNGKEPSRGI NPDEAVAYGAAVQAGVLSGDQDTGDLVLLDVCPLTLGIETVGGVMTKLIPRNTVVPTKKSQIFSTASDNQPTV TIKVYEGERPLTKDNHLLGTFDLTGIPPAPRGVPQIEVTFEIDVNGILRVTAEDKGTGNKNKITITNDQNRLTPEEI ERMVNDAEKFAEEDKCLKERIDTRNELESYAYSLKNQIGDKELGGKLSSEDKETMEKAVEEKIEWLESHQDA DIEDFKAKKKELEEIVQPIISKLYGSGGPPPTGEEDTSEKDEL

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Tris-based buffer, 50% glycerol

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months.

Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Endoplasmic reticulum chaperone that plays a key role in protein folding and quality control in the endoplasmic reticulum lumen. Involved in the correct folding of proteins and degradation of misfolded proteins via its interaction with DNAJC10/ERdj5, probably to facilitate the release of DNAJC10/ERdj5 from its substrate. Acts as a key repressor of the ERN1/IRE1-mediated unfolded protein response (UPR). In the unstressed endoplasmic reticulum, recruited by DNAJB9/ERdj4 to the luminal region of ERN1/IRE1, leading to disrupt the dimerization of ERN1/IRE1, thereby inactivating ERN1/IRE1. Accumulation of misfolded protein in the endoplasmic reticulum causes release of HSPA5/BiP from ERN1/IRE1, allowing homodimerization and subsequent activation of ERN1/IRE1. Plays an auxiliary role in post-translational transport of small presecretory proteins across endoplasmic reticulum (ER). May function as an allosteric modulator for SEC61 channel-forming translocon complex, likely cooperating with SEC62 to enable the productive insertion of these precursors into SEC61 channel. Appears to specifically regulate translocation of precursors having inhibitory residues in their mature region that weaken channel gating. May also play a role in apoptosis and cell proliferation.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481