

## RBBP9 Protein, Mouse, Recombinant (His)

## General Information

Synonyms:	Rbbp9;B5T-overexpressed gene protein (Protein BOG);Putative hydrolase RBBP9;Bog; Retinoblastoma-binding protein 9 (RBBP-9)
Protein Construction:	1-186 aa
Species:	Mouse
Expression Host:	P. pastoris (Yeast)
Accession:	O88851
Molecular Weight:	22.4 kDa (predicted)
AA Sequence:	MASPNKAVIVPGNGGGDVATHGWYGVKKGLEQIPGFQCLAKNMPDPITARESILWLPFMETELHCDEKTIIG HSSGAIAAMRYAETHQVYALVLSAYTSDLGDENERASGYFSRPWQWEKIKANCPHIVQFGSTDDPFLPWKE QQEVADRLDAKLYKFTDRGHFQNTTEFHILISVVKSMKGP

## QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

## Preparation and Storage

## Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

## Stability &amp; Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

## Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

## Protein Background

Serine hydrolase whose substrates have not been identified yet. May negatively regulate basal or autocrine TGF-beta signaling by suppressing SMAD2-SMAD3 phosphorylation. May play a role in the transformation process due

## A DRUG SCREENING EXPERT

---

to its capacity to confer resistance to the growth-inhibitory effects of TGF-beta through interaction with RB1 and the subsequent displacement of E2F1.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481