

## Ag85C Protein, Mycobacterium tuberculosis, Recombinant (His)

## General Information

Synonyms:	Antigen 85 complex C (85C;Ag85C);Acyl-CoA:diacylglycerol acyltransferase;Diacylglycerol acyltransferase/mycolyltransferase Ag85C;fbpC;DGAT;mpt45;Fibronectin-binding protein C (Fbps C)
Protein Construction:	46-340 aa
Species:	Mycobacterium tuberculosis
Expression Host:	E. coli
Accession:	P9WQN8
Molecular Weight:	36.1 kDa (predicted)
AA Sequence:	AFSRPGLPVEYLQVPSASMGRDIKVQFQGGGPHAVYLLDGLRAQDDYNGWDINTPAFEEYYQSGLSVIMPV GGQSSFYTDWYQPSQSNGQNYTYKWETFLTREMPAWLQANKGVSPTGNAAVGLSMSGGALILAAAYYPQ QFPYAASLSGFLNPSEGWWPTLIGLAMNDSGGYNANSMWGPSSDPAWKRNDPMVQIPRLVANNTRIWVY CGNGTPSDLGGDNIPAKFLEGLTLRTNQTFRDTYAADGGRNGVFNFPNGTHSWPYWNEQLVAMKADIQH VLNGATPPAAPAAPAA

## QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Tris-based buffer, 50% glycerol

## Preparation and Storage

## Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

## Stability &amp; Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

## Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

## Protein Background

The antigen 85 proteins (FbpA, FbpB, FbpC) are responsible for the high affinity of mycobacteria to fibronectin, a

## A DRUG SCREENING EXPERT

---

large adhesive glycoprotein, which facilitates the attachment of M.tuberculosis to murine alveolar macrophages (AMs). They also help to maintain the integrity of the cell wall by catalyzing the transfer of mycolic acids to cell wall arabinogalactan and through the synthesis of alpha,alpha-trehalose dimycolate (TDM, cord factor). They catalyze the transfer of a mycoloyl residue from one molecule of alpha,alpha-trehalose monomycolate (TMM) to another TMM, leading to the formation of TDM.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481