

Cytotoxin 2 Protein, Naja atra, Recombinant (His & Myc)

General Information

Synonyms:	Cytotoxin 2;Cardiotoxin 1A;Cardiotoxin analog II;Cardiotoxin 2 (CTX-2);Cardiotoxin II; Cardiotoxin A2 (CTX A2)
Protein Construction:	22-81 aa
Species:	Naja atra
Expression Host:	E. coli
Accession:	P01442
Molecular Weight:	14.2 kDa (predicted)
AA Sequence:	LKCNKLVPLFYKTCPAGKNLCYKMFVSNLTVPVKRGCIDVCPKNSALVKYVCCNTDRCN

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 85% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Basic protein that binds to cell membrane and depolarizes cardiomyocytes. It also shows lytic activities, but 2-fold less important than that of CTX-A4. It binds to the integrin alpha-V/beta-3 (ITGAV/ITGB3) with a moderate affinity. It may interact with sulfatides in the cell membrane which induces pore formation and cell internalization and is

A DRUG SCREENING EXPERT

responsible for cytotoxicity in cardiomyocytes. It also may target the mitochondrial membrane and induce mitochondrial swelling and fragmentation.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481