

IL-5 Protein, Pig, Recombinant (His)

General Information

Synonyms:	IL-5; Interleukin-5; Eosinophil differentiation factor; T-cell replacing factor (TRF); IL5
Protein Construction:	22-134 aa
Species:	Sus scrofa (Pig)
Expression Host:	P. pastoris (Yeast)
Accession:	Q9MYM5
Molecular Weight:	15.0 kDa (predicted)
AA Sequence:	VENTMNRLLVAETLTLSSIHRTLLIGDGNLMISTPVHTNHQLCIEEVFQGITLKNQTARGDAVEKLFQNLSLIKE YIDRQKKNCGGERWRVTQFLDYLQVFLGVINTEWTMES

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 85% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Tris-based buffer, 50% glycerol

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Factor that induces terminal differentiation of late-developing B-cells to immunoglobulin secreting cells.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481