

Angiotensin-2 Protein, Rat, Recombinant (His)

General Information

Synonyms: Angiotensin-2;Angpt2;Agpt2;ANG-2

Protein Construction: 19-496 aa

Species: Rat

Expression Host: Baculovirus Insect Cells

Accession: O35462

Molecular Weight: 60.1 kDa (predicted)

AA Sequence: YNNFRKSVSDSTGRRQYQVQNGPCSYTFLLPETDSCRSSSSPYMSNAVQRDAPLDYDDSVQRLQVLENILEN
NTQWLMKLENYIQDNMKKEMVEIQQNVVQNQTAVMIEIGTSLNQTAAQTRKLTQVLAQVNLQVTRLELQL
LQHSISTNKLEKQILDQTSEINKLQDKNSFLEKKVLDMEDKHSVQLQSMKEQKQDQLQVLVSKQSSVIDELEKK
LVTATVNNSVLQKQHDLMETVNSLLTMMSSPDYKSSVAVPKKEKTTFRDCAEIFKSLTTSGIYTLTFPNSTE
EVKAYCDMDMGGGGWTVIQHREDGSDVFQRTWKEYKEGFGSPLGEYWLGNFVSQLTSGHRYVLKIQKLD
WEGSEAHSLYEHFYLSGEESNYRIHLTGLTGTAGKISSISQPGSDFSTKSDNDKCICKCSQMLTGGWWFDAC
GPSNLNGQYYPQKQNTNKFNGIKWYYWKGGSGYSLKATTMMIRPADF

QC Testing

Biological Activity: Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.

Purity: > 85% as determined by SDS-PAGE.

Endotoxin: < 1.0 EU/ μ g of the protein as determined by the LAL method.

Formulation: If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μ g/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Binds to TEK/TIE2, competing for the ANGPT1 binding site, and modulating ANGPT1 signaling. Can induce tyrosine phosphorylation of TEK/TIE2 in the absence of ANGPT1. In the absence of angiogenic inducers, such as VEGF, ANGPT2-mediated loosening of cell-matrix contacts may induce endothelial cell apoptosis with consequent vascular regression. In concert with VEGF, it may facilitate endothelial cell migration and proliferation, thus serving as a permissive angiogenic signal.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481