

MBL2 Protein, Rat, Recombinant (His)

General Information

Synonyms:	Ra-reactive factor polysaccharide-binding component p28A (RaRF p28A);Mannan-binding protein;Mannose-binding protein C;MBP-C;Mbl2
Protein Construction:	19-244 aa
Species:	Rat
Expression Host:	E. coli
Accession:	P08661
Molecular Weight:	30.0 kDa (predicted)
AA Sequence:	ETLTEGAQSSCPVIACSSPGLNGFPGKDGHDGAKGEKGEPEGQLRGLQGPPGKVGPPAGPPGNPGSKGATGPKGDRGESVEFDTTNIDLEIAALRSELRAMRKWVLLSMSENVGKKYFMSSVRRMPLNRAKALCSELQGTVATPRNAEENRAIQNVAKDVAFLGITDQRTENVFEDLTGNRVRYTNWNEGEPNNVSGGENCVLLTNGKWNDVPCSDSFLVCFSD

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

A DRUG SCREENING EXPERT

Calcium-dependent lectin involved in innate immune defense. Binds mannose, fucose and N-acetylglucosamine on different microorganisms and activates the lectin complement pathway. Binds to late apoptotic cells, as well as to apoptotic blebs and to necrotic cells, but not to early apoptotic cells, facilitating their uptake by macrophages.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481