

ECHS1 Protein, Mouse, Recombinant (HA & His)

General Information

Synonyms:	Enoyl-CoA hydratase, mitochondrial;Short-chain enoyl-CoA hydratase (SCEH);mECH;Enoyl-CoA hydratase 1 (ECHS1);mECH1;mitochondrial;Echs1
Protein Construction:	28-290 aa (C225S)
Species:	Mouse
Expression Host:	E. coli
Accession:	Q8BH95
Molecular Weight:	35.6 kDa (predicted)
AA Sequence:	ASGANFQYIITEKKGKNSVGLIQLNRPKALNALCNGLIEELNQALETFEQDPAVGAIVLTTGGDKAFAAGADIK EMQNRTFQDCYSSKFLSHWDHITRVKPKVIAAVNGYALGGGCELAMMCDIYAGEKAQFGQPEILLGTIPGAG GTQRLTRAVGKSLAMEMVLTDGRISAQDAKQAGLVSKIFPVEKLVEEAIQSAEKIASNSKIVVAMAKESVNAA FEMTLTEGNKLEKRLFYSTFATDDRREGMTAFVEKRKANFKDH

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 85% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

A DRUG SCREENING EXPERT

Straight-chain enoyl-CoA thioesters from C4 up to at least C16 are processed, although with decreasing catalytic rate. Has high substrate specificity for crotonyl-CoA and moderate specificity for acryloyl-CoA, 3-methylcrotonyl-CoA and methacrylyl-CoA. It is noteworthy that binds tiglyl-CoA, but hydrates only a small amount of this substrate.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481