


## Pseudoginsenoside F11

## Chemical Properties

CAS No. :	69884-00-0
Formula:	C42H72O14
Molecular Weight:	801.01
Storage:	Powder: -20°C for 3 years   In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



## Biological Description

Description	Pseudoginsenoside F11 (Ginsenoside A1) is a component of Panax quinquefolium (American ginseng) which main pharmacological activities are positive inotropic effect on isolated heart function, good therapeutic effect on myocardial ischemia, and good protective effect on cardio-cerebrovascular system and nervous system.
Targets(IC50)	Endogenous Metabolite

## Solubility Information

Solubility	DMSO: 50 mg/mL (62.42 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (2.5 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

## Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.2484 mL	6.2421 mL	12.4842 mL
5 mM	0.2497 mL	1.2484 mL	2.4968 mL
10 mM	0.1248 mL	0.6242 mL	1.2484 mL
50 mM	0.025 mL	0.1248 mL	0.2497 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

Wang H, et al. The pseudoginsenoside F11 ameliorates cisplatin-induced nephrotoxicity without compromising its anti-tumor activity in vivo. *Scientific Reports* [2014, 4:41986]

Zhu L, et al. Pseudoginsenoside-F11 attenuates cognitive dysfunction and tau phosphorylation in sporadic Alzheimer's disease rat model. *Acta Pharmacol Sin.* 2021 Sep;42(9):1401-1408.

Fu X, et al. Pseudoginsenoside F11 ameliorates the dysfunction of the autophagy-lysosomal pathway by activating calcineurin-mediated TFEB nuclear translocation in neuron during permanent cerebral ischemia. *Exp Neurol.* 2021 Apr;338:113598.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481