

HLA-A*02:01&B2M&LMP2 (CLGGLTMV) Tetramer Protein, Human, MHC (His & Avi)

General Information

Synonyms:	RMF;PSMB9;Proteasome chain 7;RING12;LMP2;LMP-2;MHC;Macropain chain 7
Protein Construction:	Gly25-Thr305(HLA-A*02:01),Ile21-Met119(B2M) and CLGGLTMV peptide. Tetramer is assembled by biotinylated monomer and streptavidin.
Species:	Human
Expression Host:	HEK293 Cells
Accession:	A0A140T913(HLA-A*02:01)&P61769(B2M)&CLGGLTMV
Molecular Weight:	The protein has a predicted MW of 258 kDa. Due to glycosylation, the protein migrates to 260-265 kDa under Non reducing (N) condition based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Immobilized Human HLA-A*02:01&B2M&LMP2 (CLGGLTMV) Tetramer, His Tag at 2µg/ml (100µl/Well) on the plate. Dose response curve for Anti-HLA-A*02:01&B2M&LMP2 (CLGGLTMV) Antibody, hFc Tag with the EC50 of 9.6ng/ml determined by ELISA.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 µg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months.

Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

The immunoproteasome, having been linked to neurodegenerative diseases and hematological cancers, has been shown to play an important role in MHC class I antigen presentation. The development of molecular probes that selectively inhibit the major catalytic subunit, LMP2, of the immunoproteasome, LMP2-rich cancer cells compared to LMP2-deficient cancer cells are more sensitive to growth inhibition by the LMP2-specific inhibitor, implicating an

important role of LMP2 in regulating cell growth of malignant tumors that highly express LMP2.

Reference

Ho YK, et al. LMP2-specific inhibitors: chemical genetic tools for proteasome biology. Chem Biol. 2007 Apr;14(4): 419-30. doi: 10.1016/j.chembiol.2007.03.008. PMID: 17462577; PMCID: PMC5541682.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481