

Picoside I


Chemical Properties

CAS No. : 27409-30-9

Formula: C₂₄H₂₈O₁₁

Molecular Weight: 492.47

Storage: Store at low temperature, Keep away from direct sunlight
 Powder: -20°C for 3 years | In solvent: -80°C for 1 year
 Actual storage temperature shall be subject to the COA.



Biological Description

Description	Picoside I (6'-Cinnamoylcatalpol), a hepatoprotective agent, is reported to be antimicrobial and used against hepatitis B. It has antioxidant, and anti-inflammatory activities, it may be the valuable anti-invasive drug candidates for cancer therapy by suppressing Collagenases and Gelatinases. Picoside I can enhance basic fibroblast growth factor(bFGF)-, staurosporine- or dbc-mitogen-activated protein (MAP)-induced neurite outgrowth from PC12D cells.
Targets(IC50)	MMP,STAT
In vitro	Picoside I and Picoside II caused a concentration-dependent (> 0.1 microM) enhancement of basic fibroblast growth factor (bFGF, 2 ng/ml)-, staurosporine (10 nM)- and dibutyryl cyclic AMP (dbcAMP, 0.3 mM)-induced neurite outgrowth from PC12D cells. Furthermore, picosides-induced enhancements of the bFGF-action were markedly inhibited by GF109203X (0.1 microM), a protein kinase C inhibitor. The expression of phosphorylated MAP kinase was markedly increased by bFGF (2 ng/ml) and dbcAMP (0.3 mM), whereas that was not enhanced by staurosporine (10 nM). Picosides had no effect on the phosphorylation of MAP kinase induced by bFGF or dbcAMP and also unaffected it in the presence of staurosporine. These results suggest that Picoside I and Picoside II enhance bFGF-, staurosporine- or dbcAMP-induced neurite outgrowth from PC12D cells, probably by amplifying a down-stream step of MAP kinase in the intracellular MAP kinase-dependent signaling pathway.

Solubility Information

Solubility	H ₂ O: 10 mg/mL (20.31 mM),Sonication is recommended. Ethanol: 10 mg/mL (20.31 mM),Sonication is recommended. DMSO: 55 mg/mL (111.68 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.0306 mL	10.1529 mL	20.3058 mL
5 mM	0.4061 mL	2.0306 mL	4.0612 mL
10 mM	0.2031 mL	1.0153 mL	2.0306 mL
50 mM	0.0406 mL	0.2031 mL	0.4061 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Chander R, et al. Picroliv, picroside-I and kutkoside from Picrorhiza kurrooa are scavengers of superoxide anions. *Biochem Pharmacol.* 1992 Jul 7;44(1):180-3.

Huang B, Lin B, Zheng H, et al. Discovery of natural products as influenza neuraminidase inhibitors: in silico screening, in vitro validation, and molecular dynamic simulation studies. *Molecular Diversity.* 2025: 1-17.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481