

Anti-TGFBR2 Monoclonal Antibody

Product Details

Ig Type:	Rabbit monoclonal IgG
Reactivity:	Human
Conjugation:	Unconjugated
Molecular Weight:	150 kDa
Purification:	Protein A Affinity Purified

Applications

Verified Activity:	Flow cytometry analysis of TGFBR2 overexpressed 311F cells with TMAZ-0192, followed by goat anti-rabbit IgG-ABflo 647 (red line). The isotype control is rabbit IgG (black line).
Application:	ELISA, FCM, IF
Recommended	0.1-0.2 µg/10E6 cells for FCM; 1 ng/µl for ELISA

Properties

Purity:	> 95% as determined by SDS-PAGE.
Stability & Storage:	Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	TGFBR2
Antigen Species:	Human
Gene ID:	7048
Uniprot ID:	P37173
Synonyms:	Transforming growth factor-beta receptor type II;TGF-beta type II receptor;TGF-beta receptor type-2
Biology Area:	Drug Target

Research Background

Transmembrane serine/threonine kinase forming with the TGF-beta type I serine/threonine kinase receptor, TGFBR1, the non-promiscuous receptor for the TGF-beta cytokines TGFB1, TGFB2 and TGFB3. Transduces the TGFB1, TGFB2 and TGFB3 signal from the cell surface to the cytoplasm and thus regulates a plethora of physiological and pathological processes including cell cycle arrest in epithelial and hematopoietic cells, control of mesenchymal cell proliferation and differentiation, wound healing, extracellular matrix production, immunosuppression and carcinogenesis. The formation of the receptor complex composed of 2 TGFBR1 and 2 TGFBR2 molecules symmetrically bound to the cytokine dimer results in the phosphorylation and activation of TGFBR1 by the constitutively active TGFBR2. Activated TGFBR1 phosphorylates SMAD2 which dissociates from the receptor and interacts with SMAD4. The SMAD2-SMAD4 complex is subsequently translocated to the nucleus where it modulates the transcription of the TGF-beta-regulated genes. This constitutes the canonical SMAD-dependent TGF-beta signaling cascade. Also involved in non-canonical, SMAD-independent TGF-beta signaling pathways|Has transforming growth factor beta-activated receptor activity|Has transforming growth factor beta-activated receptor

A DRUG SCREENING EXPERT

activity|Binds TGFB1, TGFB2 and TGFB3 in the picomolar affinity range without the participation of additional receptors. Blocks activation of SMAD2 and SMAD3 by TGFB1

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481