

Anti-PD-L1 Monoclonal Antibody-Biotin

Product Details

Ig Type:	Rabbit monoclonal IgG
Reactivity:	Human
Conjugation:	Biotin
Molecular Weight:	150 kDa
Purification:	Protein A Affinity Purified

Applications

Verified Activity:	Flow cytometry analysis of PD-L1 overexpressed 293F cells with TMAZ-0055BBI, which was counjugated with SA-PE (red). The WT 293F control is black line.
Application:	ELISA, FACS, IF
Recommended	0.1-0.2 µg/10E6 cells for FACS; 1 ng/µl for ELISA

Properties

Purity:	Greater than 95%, as determined by SDS-PAGE.
Stability & Storage:	For unconjugated and biotin antibody, store at -20°C or -80°C for long term. Avoid repeated freeze-thaw cycles. For APC/PE/AF647/AF555/AF488 conjugated antibody, store at 4°C, do not freeze.
Shipping:	For unconjugated and biotin antibody, shipping with dry ice. For APC/PE/AF647/AF555/AF488 conjugated antibody, shipping with blue ice.

Antigen Details

Immunogen:	PD-L1
Antigen Species:	Human
Gene ID:	29126
Uniprot ID:	Q9NZQ7
Synonyms:	B7 homolog 1;Programmed cell death 1 ligand 1
Biology Area:	Immunology Research

Research Background

Plays a critical role in induction and maintenance of immune tolerance to self (PubMed: 11015443, PubMed: 28813410, PubMed: 28813417, PubMed: 31399419). As a ligand for the inhibitory receptor PDCD1/PD-1, modulates the activation threshold of T-cells and limits T-cell effector response (PubMed: 11015443, PubMed: 28813410, PubMed: 28813417, PubMed: 36727298). Through a yet unknown activating receptor, may costimulate T-cell subsets that predominantly produce interleukin-10 (IL10) (PubMed: 10581077). Can also act as a transcription coactivator: in response to hypoxia, translocates into the nucleus via its interaction with phosphorylated STAT3 and promotes transcription of GSDMC, leading to pyroptosis (PubMed: 32929201)|The PDCD1-mediated inhibitory pathway is exploited by tumors to attenuate anti-tumor immunity and escape destruction by the immune system, thereby facilitating tumor survival (PubMed: 28813410, PubMed: 28813417). The interaction with PDCD1/PD-1 inhibits cytotoxic T lymphocytes (CTLs) effector function (By similarity). The blockage of the PDCD1-mediated pathway results

A DRUG SCREENING EXPERT

in the reversal of the exhausted T-cell phenotype and the normalization of the anti-tumor response, providing a rationale for cancer immunotherapy (By similarity)

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481


Anti-PD-L1 Monoclonal Antibody-Biotin

Product Details

Ig Type:	Rabbit monoclonal IgG
Reactivity:	Human
Conjugation:	Biotin
Molecular Weight:	150 kDa
Clone:	A1AA001
Purification:	Protein A Affinity Purified

Applications

Verified Activity: Flow cytometry analysis of PD-L1 overexpressed 293F cells with TMAZ-0055BBI, which was conjugated with SA-PE (red). The WT 293F control is black line.



Application:	ELISA, FACS, IF
Recommended	0.1-0.2 µg/10E6 cells for FACS; 1 ng/µl for ELISA

Properties

Purity:	Greater than 95%, as determined by SDS-PAGE.
Stability & Storage:	For unconjugated and biotin antibody, store at -20°C or -80°C for long term. Avoid repeated freeze-thaw cycles. For APC/PE/AF647/AF555/AF488 conjugated antibody, store at 4°C, do not freeze.
Shipping:	For unconjugated and biotin antibody, shipping with dry ice. For APC/PE/AF647/AF555/AF488 conjugated antibody, shipping with blue ice.

Antigen Details

Immunogen:	PD-L1
Antigen Species:	Human
Gene ID:	29126
Uniprot ID:	Q9NZQ7
Synonyms:	B7 homolog 1; Programmed cell death 1 ligand 1
Biology Area:	Immunology Research
Classification:	Antigen-presenting cells, Dendritic cells, Monocytes

Research Background

Plays a critical role in induction and maintenance of immune tolerance to self (PubMed: 11015443, PubMed: 28813410, PubMed: 28813417, PubMed: 31399419). As a ligand for the inhibitory receptor PDCD1/PD-1, modulates the activation threshold of T-cells and limits T-cell effector response (PubMed: 11015443, PubMed: 28813410, PubMed: 28813417, PubMed: 36727298). Through a yet unknown activating receptor, may costimulate T-cell subsets that predominantly produce interleukin-10 (IL10) (PubMed: 10581077). Can also act as a transcription coactivator: in response to hypoxia, translocates into the nucleus via its interaction with phosphorylated STAT3 and promotes transcription of GSDMC, leading to pyroptosis (PubMed: 32929201)|The PDCD1-mediated inhibitory pathway is exploited by tumors to attenuate anti-tumor immunity and escape destruction by the immune system, thereby facilitating tumor survival (PubMed: 28813410, PubMed: 28813417). The interaction with PDCD1/PD-1 inhibits cytotoxic T lymphocytes (CTLs) effector function (By similarity). The blockage of the PDCD1-mediated pathway results in the reversal of the exhausted T-cell phenotype and the normalization of the anti-tumor response, providing a rationale for cancer immunotherapy (By similarity)

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481

Anti-PD-L1 Monoclonal Antibody-Biotin

Product Details

Ig Type:	Rabbit monoclonal IgG
Reactivity:	Human
Conjugation:	Biotin
Molecular Weight:	150 kDa
Purification:	Protein A Affinity Purified

Applications

Verified Activity:	Flow cytometry analysis of PD-L1 overexpressed 293F cells with TMAZ-0055BBI, which was counjugated with SA-PE (red). The WT 293F control is black line.
Application:	ELISA, FACS, IF
Recommended	0.1-0.2 µg/10E6 cells for FACS; 1 ng/µl for ELISA

Properties

Purity:	Greater than 95%, as determined by SDS-PAGE.
Stability & Storage:	For unconjugated and biotin antibody, store at -20°C or -80°C for long term. Avoid repeated freeze-thaw cycles. For APC/PE/AF647/AF555/AF488 conjugated antibody, store at 4°C, do not freeze.
Shipping:	For unconjugated and biotin antibody, shipping with dry ice. For APC/PE/AF647/AF555/AF488 conjugated antibody, shipping with blue ice.

Antigen Details

Immunogen:	PD-L1
Antigen Species:	Human
Gene ID:	29126
Uniprot ID:	Q9NZQ7
Synonyms:	B7 homolog 1;Programmed cell death 1 ligand 1
Biology Area:	Immunology Research

Research Background

Plays a critical role in induction and maintenance of immune tolerance to self (PubMed: 11015443, PubMed: 28813410, PubMed: 28813417, PubMed: 31399419). As a ligand for the inhibitory receptor PDCD1/PD-1, modulates the activation threshold of T-cells and limits T-cell effector response (PubMed: 11015443, PubMed: 28813410, PubMed: 28813417, PubMed: 36727298). Through a yet unknown activating receptor, may costimulate T-cell subsets that predominantly produce interleukin-10 (IL10) (PubMed: 10581077). Can also act as a transcription coactivator: in response to hypoxia, translocates into the nucleus via its interaction with phosphorylated STAT3 and promotes transcription of GSDMC, leading to pyroptosis (PubMed: 32929201)|The PDCD1-mediated inhibitory pathway is exploited by tumors to attenuate anti-tumor immunity and escape destruction by the immune system, thereby facilitating tumor survival (PubMed: 28813410, PubMed: 28813417). The interaction with PDCD1/PD-1 inhibits cytotoxic T lymphocytes (CTLs) effector function (By similarity). The blockage of the PDCD1-mediated pathway results

A DRUG SCREENING EXPERT

in the reversal of the exhausted T-cell phenotype and the normalization of the anti-tumor response, providing a rationale for cancer immunotherapy (By similarity)

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481


Anti-PD-L1 Monoclonal Antibody-Biotin

Product Details

Ig Type:	Rabbit monoclonal IgG
Reactivity:	Human
Conjugation:	Biotin
Molecular Weight:	150 kDa
Clone:	A1AA001
Purification:	Protein A Affinity Purified

Applications

Verified Activity: Flow cytometry analysis of PD-L1 overexpressed 293F cells with TMAZ-0055BBI, which was counjugated with SA-PE (red). The WT 293F control is black line.



Application:	ELISA, FCM, IF
Recommended	0.1-0.2 µg/10E6 cells for FCM; 1 ng/µl for ELISA

Properties

Purity:	> 95% as determined by SDS-PAGE.
Stability & Storage:	Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	PD-L1
Antigen Species:	Human
Gene ID:	29126
Uniprot ID:	Q9NZQ7
Synonyms:	B7 homolog 1; Programmed cell death 1 ligand 1
Biology Area:	Immunology Research

Research Background

Plays a critical role in induction and maintenance of immune tolerance to self (PubMed: 11015443, PubMed: 28813410, PubMed: 28813417, PubMed: 31399419). As a ligand for the inhibitory receptor PDCD1/PD-1, modulates the activation threshold of T-cells and limits T-cell effector response (PubMed: 11015443, PubMed: 28813410,

A DRUG SCREENING EXPERT

PubMed: 28813417, PubMed: 36727298). Through a yet unknown activating receptor, may costimulate T-cell subsets that predominantly produce interleukin-10 (IL10) (PubMed: 10581077). Can also act as a transcription coactivator: in response to hypoxia, translocates into the nucleus via its interaction with phosphorylated STAT3 and promotes transcription of GSDMC, leading to pyroptosis (PubMed: 32929201)|The PDCD1-mediated inhibitory pathway is exploited by tumors to attenuate anti-tumor immunity and escape destruction by the immune system, thereby facilitating tumor survival (PubMed: 28813410, PubMed: 28813417). The interaction with PDCD1/PD-1 inhibits cytotoxic T lymphocytes (CTLs) effector function (By similarity). The blockage of the PDCD1-mediated pathway results in the reversal of the exhausted T-cell phenotype and the normalization of the anti-tumor response, providing a rationale for cancer immunotherapy (By similarity)

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481


Anti-PD-L1 Monoclonal Antibody-Biotin

Product Details

Ig Type:	Rabbit monoclonal IgG
Reactivity:	Human
Conjugation:	Biotin
Molecular Weight:	150 kDa
Clone:	A1AA001
Purification:	Protein A Affinity Purified

Applications

Verified Activity: Flow cytometry analysis of PD-L1 overexpressed 293F cells with TMAZ-0055BBI, which was conjugated with SA-PE (red). The WT 293F control is black line.



Application:	ELISA, FACS, IF
Recommended	0.1-0.2 µg/10E6 cells for FACS; 1 ng/µl for ELISA

Properties

Purity:	Greater than 95%, as determined by SDS-PAGE.
Stability & Storage:	For unconjugated and biotin antibody, store at -20°C or -80°C for long term. Avoid repeated freeze-thaw cycles. For APC/PE/AF647/AF555/AF488 conjugated antibody, store at 4°C, do not freeze.
Shipping:	For unconjugated and biotin antibody, shipping with dry ice. For APC/PE/AF647/AF555/AF488 conjugated antibody, shipping with blue ice.

Antigen Details

Immunogen:	PD-L1
Antigen Species:	Human
Gene ID:	29126
Uniprot ID:	Q9NZQ7
Synonyms:	B7 homolog 1; Programmed cell death 1 ligand 1
Biology Area:	Immunology Research
Classification:	Antigen-presenting cells, Dendritic cells, Monocytes

Research Background

Plays a critical role in induction and maintenance of immune tolerance to self (PubMed: 11015443, PubMed: 28813410, PubMed: 28813417, PubMed: 31399419). As a ligand for the inhibitory receptor PDCD1/PD-1, modulates the activation threshold of T-cells and limits T-cell effector response (PubMed: 11015443, PubMed: 28813410, PubMed: 28813417, PubMed: 36727298). Through a yet unknown activating receptor, may costimulate T-cell subsets that predominantly produce interleukin-10 (IL10) (PubMed: 10581077). Can also act as a transcription coactivator: in response to hypoxia, translocates into the nucleus via its interaction with phosphorylated STAT3 and promotes transcription of GSDMC, leading to pyroptosis (PubMed: 32929201)|The PDCD1-mediated inhibitory pathway is exploited by tumors to attenuate anti-tumor immunity and escape destruction by the immune system, thereby facilitating tumor survival (PubMed: 28813410, PubMed: 28813417). The interaction with PDCD1/PD-1 inhibits cytotoxic T lymphocytes (CTLs) effector function (By similarity). The blockage of the PDCD1-mediated pathway results in the reversal of the exhausted T-cell phenotype and the normalization of the anti-tumor response, providing a rationale for cancer immunotherapy (By similarity)

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481