


Aknadinine

Chemical Properties

CAS No. :	24148-86-5
Formula:	C ₂₀ H ₂₅ N ₁ O ₅
Molecular Weight:	359.42
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	Aknadinine is a biochemical isolated from <i>Stephania sutchuenensis</i> .
Targets(IC50)	Others

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.7823 mL	13.9113 mL	27.8226 mL
5 mM	0.5565 mL	2.7823 mL	5.5645 mL
10 mM	0.2782 mL	1.3911 mL	2.7823 mL
50 mM	0.0556 mL	0.2782 mL	0.5565 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

- He L, Zhang YH, Tang LJ, Song SH, Sun QY. [Study on the alkaloids in the stems and leaves of *Stephania cepharantha* (II)]. *Zhong Yao Cai*. 2010 Oct;33(10):1568-70. Chinese. PubMed PMID: 21355192.
- Tang L, Zhang Y, Hao X, He L, Song S, Hao X, Yang X. [Hasubanan type alkaloids in *Stephania hernandifolia*]. *Zhongguo Zhong Yao Za Zhi*. 2010 Aug;35(15):1973-7. Chinese. PubMed PMID: 20931849.
- Nielsen DK, Nielsen LL, Jones SB, Toll L, Asplund MC, Castle SL. Synthesis of isohasubanan alkaloids via enantioselective ketone allylation and discovery of an unexpected rearrangement. *J Org Chem*. 2009 Feb 6;74(3):1187-99. doi: 10.1021/jo802370v. PubMed PMID: 19072324.
- Liu WK, Wang XK, Che CT. Cytotoxic effects of sinococuline. *Cancer Lett*. 1996 Feb 6;99(2):217-24. PubMed PMID: 8616827.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481