

## Fusidic acid

## Chemical Properties

CAS No. :	6990-06-3
Formula:	C <sub>31</sub> H <sub>48</sub> O <sub>6</sub>
Molecular Weight:	516.71
Storage:	Powder: -20°C for 3 years   In solvent: -80°C for 1 year <small>Actual storage temperature shall be subject to the COA.</small>

## Biological Description

Description	Fusidic acid (Fusidine) is an antibiotic isolated from the fermentation broth of <i>Fusidium coccineum</i> .
Targets(IC50)	Antibacterial, Antibiotic
Kinase Assay	DHODase activity is measured by the DCIP colorimetric assay. This is a coupled assay in which oxidation of DHO and subsequent reduction of ubiquinone are stoichiometrically equivalent to the reduction of DCIP. Reduction of DCIP is accompanied by a loss of absorbance at 610 nm ( $\epsilon=21500$ M/cm). The assay is performed in a 96-well microtiter plate at ambient temperature (ca. 25°C). Stock solutions of 10 mM leflunomide and A771726 are prepared in dimethyl sulfoxide (DMSO) and these are diluted with reaction buffer (100 mM Tris and 0.1 % Triton X-100, pH 8.0) to prepare working stocks of the inhibitors at varying concentrations. For each reaction, the well contained 10 nM DHODase, 68 $\mu$ M DCIP, 0.16 mg/mL gelatin, the stated concentration of ubiquinone, 10 $\mu$ L of an inhibitor working stock to give the stated final concentration, and reaction buffer. After a 5-min equilibration period, the reaction is initiated by addition of DHO to the stated final concentrations. The total volume of reaction mixture for each assay is 150 $\mu$ L, and the final DMSO concentration is $\leq 0.01\%$ (v/v). The reaction progress is followed by recording the loss of absorbance at 610 nm over a 10-min period (during which the velocity remained linear). Velocities are reported as the change in absorbance at 610 nm per minute, and each reported value is the average of three replicates. In experiments where the DHO or ubiquinone concentration is varied, the other substrate is held constant at 200 $\mu$ M. To determine the inhibitor potency of leflunomide and A771726, the effects of varying concentrations of the two compounds on the initial velocity of the DHODase reaction is measured over a concentration range of 0.01-1.0 $\mu$ M. In these experiments the DHO and ubiquinone concentrations are held constant at 200 and 100 $\mu$ M, respectively.

## Solubility Information

Solubility	DMSO: 125 mg/mL (241.92 mM), Sonication is recommended. ( $< 1$ mg/ml refers to the product slightly soluble or insoluble)
------------	---

In vivo Formulation	<p>10% DMSO+90% Saline: &lt; 10 mg/mL (19.35 mM), Lower concentrations may be soluble, but exact solubility limit is unknown.</p> <p>10% DMSO+40% PEG300+5% Tween 80+45% Saline: 10 mg/mL (19.35 mM), Solution.</p> <p><i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i></p>
---------------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.9353 mL	9.6766 mL	19.3532 mL
5 mM	0.3871 mL	1.9353 mL	3.8706 mL
10 mM	0.1935 mL	0.9677 mL	1.9353 mL
50 mM	0.0387 mL	0.1935 mL	0.3871 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Day PJ, et al. Biochemistry. 1995 May 16;34(19):6416-22.  
 Zhang H, Liang B, Sang X, et al. Discovery of Potential Inhibitors of SARS-CoV-2 Main Protease by a Transfer Learning Method. Viruses. 2023, 15(4): 891.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481