



## A DRUG SCREENING EXPERT

In vivo	c-Fos expression are reduced by a behaviorally effective dose of Istradefylline, in contrast to tropicamide by which c-Fos expression in pimozide-treated rats is actually increased. [8]
Cell Research	A CHO cell line permanently expressing the human adenosine A1 or A2A receptor is cultured in $\alpha$ -MEM supplemented with 10% (v/v) fetal bovine serum, 50 U/mL penicillin, and 50 $\mu$ g/mL streptomycin. Cells are grown at 37°C in an environment of 5% CO <sub>2</sub> . These cells are seeded on black 96-well assay plates at a density of 15,000 cells/well, and then they are cultured for 24 h.

### Solubility Information

Solubility	DMSO: 13.38 mg/mL (34.8 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

### Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.6013 mL	13.0063 mL	26.0125 mL
5 mM	0.5203 mL	2.6013 mL	5.2025 mL
10 mM	0.2601 mL	1.3006 mL	2.6013 mL
50 mM	0.052 mL	0.2601 mL	0.5203 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

- Chen JF, et al. J Neurosci, 2001, 21(10), RC143.
- Shiozaki S, et al. Psychopharmacology (Berl), 1999, 147(1), 90-95.
- Kanda T, et al. Exp Neurol, 2000, 162(2), 321-327.
- Ikeda K, et al. J Neurochem, 2002, 80(2), 262-270.
- Brambilla R, et al. Glia, 2003, 43(2), 190-194.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E\_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481