

Anti-CCL2 Monoclonal Antibody-Biotin

Product Details

Ig Type:	Rabbit monoclonal IgG
Reactivity:	Human
Conjugation:	Biotin
Molecular Weight:	150 kDa
Purification:	Protein A Affinity Purified

Applications

Verified Activity:	Flow cytometry analysis of CCL2 overexpressed 446F cells with TMAZ-0376B, followed by goat anti-rabbit IgG-ABflo 647 (red line). The isotype control is rabbit IgG (black line).
Application:	ELISA,FCM
Recommended	0.1-0.2 µg/10E6 cells for FCM; 1 ng/µl for ELISA

Properties

Purity:	> 95% as determined by SDS-PAGE.
Stability & Storage:	Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	CCL2
Antigen Species:	Human
Gene ID:	6347
Uniprot ID:	P13500
Synonyms:	HC11; Monocyte chemoattractant protein 1; C-C motif chemokine 2; Monocyte chemotactic and activating factor; Monocyte chemotactic protein 1; Monocyte secretory protein JE; Small-inducible cytokine A2
Biology Area:	Cancer Research

Research Background

Acts as a ligand for C-C chemokine receptor CCR2 (PubMed:10529171, PubMed:10587439, PubMed:9837883). Signals through binding and activation of CCR2 and induces a strong chemotactic response and mobilization of intracellular calcium ions (PubMed:10587439, PubMed:9837883). Exhibits a chemotactic activity for monocytes and basophils but not neutrophils or eosinophils (PubMed:8195247, PubMed:8627182, PubMed:9792674). May be involved in the recruitment of monocytes into the arterial wall during the disease process of atherosclerosis

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481